

Traveling Waves

1. A particular wave traveling along a string is described by:

$$y(x,t) = (0.0060\text{m})\sin\left((0.006\frac{\text{rad}}{\text{m}})x + (600\frac{\text{rad}}{\text{s}})t\right)$$

a. How much time does any particular point in the string take to move between the displacements $y=+2\text{ mm}$ and $y=-2\text{ mm}$?

Ans. There 2 interpretations to this problem.

Interpretation 1:

The period, T , for this wave is: $T = \frac{2\pi}{\omega} = 0.0105\text{ s}$

The displacement from $+2\text{mm}$ to -2mm is $\frac{1}{2}\lambda$ and therefore the time elapsed is:

$$t = \frac{T}{2} = 0.00525\text{ s}.$$


Interpretation 2:

This one is more complex. The phase difference between $+2\text{mm}$ to -2mm is:

$$\phi = \pi \left(\frac{0.0040\text{m}}{0.0120\text{m}} \right) = \frac{\pi}{3}$$

The time elapsed is related to period, T , by:

$$\frac{t}{T} = \frac{\phi}{2\pi} = \frac{\left(\frac{\pi}{3}\right)}{2\pi} \Rightarrow t = \frac{T}{6} = 0.0018\text{ s}$$



b. Determine the amplitude, frequency, period and wave speed for this traveling wave.

Ans. $y_m = 0.0060\text{ m}$, $f = \frac{\omega}{2\pi} = 95.5\text{ Hz}$, $T = \frac{1}{f} = 0.0105\text{ s}$, and $v = \frac{\omega}{k} = 1.05 \times 10^5 \frac{\text{m}}{\text{s}}$

c. What is the equation for the transverse velocity of any particular point along the string?

Ans. $v = \frac{dy}{dt} = (3.6\frac{\text{m}}{\text{s}})\sin\left[(0.006\frac{\text{rad}}{\text{m}})x + (600\frac{\text{rad}}{\text{s}})t\right]$

d. What is the maximum transverse velocity of any particular point along the string?

Ans. $v_m = \omega y_m = 3.6\text{ m/s}$

2. A sinusoidal wave of frequency 500 Hz has a speed of 350 m/s.

a. How far apart are 2 points that differ in phase by $\pi/3$ radians?

Ans. $\lambda = \frac{350\frac{\text{m}}{\text{s}}}{500\text{ Hz}} = 0.7\text{ m} \Rightarrow \frac{0.7\text{ m}}{\Delta x} = \frac{2\pi}{\left(\frac{\pi}{3}\right)} \Rightarrow \Delta x = 0.117\text{ m}$

b. What is the phase difference between 2 displacements at a certain point at times 1.000 ms apart?

$$\text{Ans. } T = \frac{1}{500 \text{ Hz}} = 0.002 \text{ s} \Rightarrow \frac{\Delta t}{T} = \frac{0.001 \text{ s}}{0.002 \text{ s}} = \frac{\phi}{2\pi} \Rightarrow \phi = \pi \text{ rad}$$

Waves on a Stretched String

3. The linear density of a string is $1.5 \times 10^{-3} \text{ kg/m}$. The equation of a transverse wave on a string is:

$$y(x,t) = (0.025 \text{ m}) \sin\left((16.0 \frac{\text{rad}}{\text{m}})x - (750 \frac{\text{rad}}{\text{s}})t\right)$$

a. What is the wave speed?

$$\text{Ans. } v = \frac{\omega}{k} = \frac{750 \frac{\text{rad}}{\text{s}}}{16.0 \frac{\text{rad}}{\text{m}}} = 46.9 \frac{\text{m}}{\text{s}}$$

b. What is the tension in the string?

$$\text{Ans. } F_T = \mu v^2 = (1.5 \times 10^{-3} \frac{\text{kg}}{\text{m}})(46.9 \frac{\text{m}}{\text{s}})^2 = 3.30 \text{ N}$$

c. What is the average power for this wave?

$$\text{Ans. } P = (\frac{1}{2})\mu v \omega^2 y_m^2 = 12.4 \text{ W}$$

4. The equation of a transverse wave on a string is:

$$y(x,t) = (2.0 \text{ mm}) \sin\left((20 \frac{\text{rad}}{\text{m}})x - (600 \frac{\text{rad}}{\text{s}})t\right)$$

The tension in the string is 15 N.

a. What is the wave speed?

$$\text{Ans. } v = \frac{\omega}{k} = \frac{600 \frac{\text{rad}}{\text{s}}}{20 \frac{\text{rad}}{\text{m}}} = 30 \frac{\text{m}}{\text{s}}$$

b. What is the linear mass density of the string, in grams per meter?

$$\text{Ans. } \mu = \frac{F_T}{v^2} = \frac{15 \text{ N}}{(30 \frac{\text{m}}{\text{s}})^2} = 1.67 \times 10^{-2} \frac{\text{kg}}{\text{m}}$$

c. Determine the average power transmitted by this wave.

$$\text{Ans. } P = (\frac{1}{2})\mu v \omega^2 y_m^2 = 0.36 \text{ W}$$

d. Demonstrate that this transverse wave satisfies the wave equation: $\frac{\partial^2 y}{\partial x^2} = \frac{1}{v^2} \frac{\partial^2 y}{\partial t^2}$.

The Wave Equation:

5. Consider the following wave: $y(x,t) = (0.05 \text{ m}) \sin\left((5.0 \frac{\text{rad}}{\text{m}})x - (10 \frac{\text{rad}}{\text{s}})t\right)$

a. Verify that this wave satisfies the wave equation: $\frac{\partial^2 y}{\partial x^2} = \frac{1}{v^2} \frac{\partial^2 y}{\partial t^2}$

$$\text{Ans. } \frac{\partial^2 y}{\partial x^2} = -(5.0 \frac{\text{rad}}{\text{m}})^2 (0.05 \text{ m}) \sin\left((5.0 \frac{\text{rad}}{\text{m}})x - (10 \frac{\text{rad}}{\text{s}})t\right) = -(1.25 \text{ m}^{-1}) \sin\left((5.0 \frac{\text{rad}}{\text{m}})x - (10 \frac{\text{rad}}{\text{s}})t\right)$$

$$\frac{1}{v^2} \frac{\partial^2 y}{\partial t^2} = -\frac{(10 \frac{\text{rad}}{\text{s}})^2 (0.05\text{m})}{(2 \frac{\text{m}}{\text{s}})^2} \sin((5.0 \frac{\text{rad}}{\text{m}})x - (10 \frac{\text{rad}}{\text{s}})t) = -(1.25\text{m}^{-1}) \sin((5.0 \frac{\text{rad}}{\text{m}})x - (10 \frac{\text{rad}}{\text{s}})t)$$

b. Use the wave equation to determine the speed of the wave.

Ans. Setting the above equations together, and not assuming a priori that $v = \omega/k$, leads to:

$$\frac{(10 \frac{\text{rad}}{\text{s}})^2 (0.05\text{m})}{v^2} = (5.0 \frac{\text{rad}}{\text{m}})^2 (0.05\text{m}) \Rightarrow v = \sqrt{v^2} = \frac{10 \frac{\text{rad}}{\text{s}}}{5.0 \frac{\text{rad}}{\text{m}}} = 2.0 \frac{\text{m}}{\text{s}}$$

c. Calculate the frequency, wavelength and period for this wave.

$$\text{Ans. } y_m = 0.05 \text{ m, } f = \frac{\omega}{2\pi} = 1.59 \text{ Hz, } \lambda = \frac{k}{2\pi} = 0.796 \text{ m, and } T = \frac{1}{f} = 0.629 \text{ s}$$

d. Consider a 2nd wave (with same frequency and wavelength) traveling in the same direction, where $y_{2\text{max}} = 0.01 \text{ m}$ and $\phi_2 = \pi/3$. Construct a phasor diagram for the combined waves.

Ans.



e. What is the amplitude of the resultant wave?

$$\text{Ans. } y' = \sqrt{y_1^2 + y_2^2 - 2y_1y_2 \cos\left(\pi - \frac{\pi}{3}\right)} = 0.0557 \text{ m}$$

Standing Waves

7. A string is attached to a string vibrator ($f=120 \text{ Hz}$) at one end and fixed at the other. The length of the string is 1.2 m .

a. What wavelengths will satisfy the standing wave condition for the string?

$$\text{Ans. } L = n\left(\frac{\lambda}{2}\right) \quad \{\text{where } n=1,2,3,\dots \text{ etc.}\}$$

b. What is the maximum wave speed that will produce a standing wave in this string?

Ans. For the same frequency, the greatest wavelength will produce the maximum speed:

$$\text{when } n=1, \lambda = \frac{2L}{n} = 2.4 \text{ m and } v = \lambda \cdot f = (2.4 \text{ m})(120 \text{ Hz}) = 288 \frac{\text{m}}{\text{s}}$$

c. What string tension will correspond to the wave in (b) is the linear density of the string is $2.5 \times 10^{-4} \text{ kg/m}$?

$$\text{Ans. } F_T = \mu v^2 = (2.5 \times 10^{-4} \frac{\text{kg}}{\text{m}})(288 \frac{\text{m}}{\text{s}})^2 = 20.7 \text{ N}$$

d. What string tension would correspond to a standing wave that produces 6 nodes in the string?

$$\text{Ans. For 6 nodes, } n=5, \lambda = \frac{2L}{n} = 0.48 \text{ m and } v = \lambda \cdot f = (0.48 \text{ m})(120 \text{ Hz}) = 57.6 \frac{\text{m}}{\text{s}}$$

$$\text{The tension in the string would be: } F_T = \mu v^2 = (2.5 \times 10^{-4} \frac{\text{kg}}{\text{m}})(57.6 \frac{\text{m}}{\text{s}})^2 = 0.829 \text{ N}$$