

**Example 4.1**

Consider the matrix  $B = \begin{bmatrix} 4 & 7 & -7 \\ 8 & -2 & 0 \\ -4 & 11 & 9 \\ -18 & -62 & 3 \\ 22 & -1 & 14 \end{bmatrix}$ .

a. What are the dimensions of the matrix?

b. What are entries  $b_{32}$ ,  $b_{23}$ ,  $b_{41}$ , and  $b_{14}$ ?

**Example 4.2**

Categorize each of the following as a vector, a column vector, or not a vector.

i.  $\begin{bmatrix} 4 \\ 0 \\ 0 \\ 7 \end{bmatrix}$

ii.  $\begin{bmatrix} 3 & -1 \\ 6 & 12 \end{bmatrix}$

iii.  $[4, 7, -18]$

iv.  $\begin{bmatrix} 5 & 0 \end{bmatrix}$

v.

**Example 4.3**

Consider the matrices  $A = \begin{bmatrix} 3 & -1 \\ 5 & 2 \\ 0 & -7 \end{bmatrix}$ ,  $B = \begin{bmatrix} 4 & 8 & -2 \\ -5 & -1 & 6 \end{bmatrix}$ , and  $C = \begin{bmatrix} -1 & -2 & 6 \\ 3 & -2 & -2 \end{bmatrix}$ .

Find  $A + B$ ,  $B - 2C$ , and  $C + A^T$

**Example 4.4**

Which of the following are column vectors?

i.  $\begin{bmatrix} 7 \\ 8 \end{bmatrix}$

ii.  $\begin{bmatrix} 3 & -1 \\ 6 & 12 \end{bmatrix}$

iii.  $[4, 7, -18]^T$

iv.  $\begin{bmatrix} 5 & 0 \end{bmatrix}$

v.

**Example 4.5**

Find  $AB$  if  $A = \begin{bmatrix} 2 & 1 \\ 4 & 6 \\ 3 & -2 \end{bmatrix}$  and  $B = \begin{bmatrix} -1 & 3 & 0 & 7 \\ -3 & 5 & -4 & -5 \end{bmatrix}$ .

**Example 4.6**

Find  $AB$  if  $A = \begin{bmatrix} 2 & 1 & 0 & 8 & -5 \\ 1 & 3 & 4 & -3 & -1 \end{bmatrix}$  and  $B = \begin{bmatrix} 3 & -2 \\ 10 & 1 \\ 2 & 4 \\ 0 & 3 \\ -6 & 7 \end{bmatrix}$ .

**Example 4.7**

Find  $AB$  if  $A = \begin{bmatrix} 2 & 1 & 0 & 8 & -5 \\ 1 & 3 & 4 & -3 & -1 \end{bmatrix}$  and  $B = \begin{bmatrix} 3 & -2 \\ 10 & 1 \\ 2 & 4 \end{bmatrix}$ .

**Example 4.8**

Write a simple dimensional proof that establishes that in general, matrix multiplication is not commutative; i.e., in general  $AB \neq BA$ .

**Example 4.9**

Use your calculator to find  $I_3 A$  and  $A I_3$  where  $A = \begin{bmatrix} 2 & 6 & -3 \\ -1 & 5 & 4 \\ 9 & 12 & -7 \end{bmatrix}$  and  $I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

**Example 4.10**


Use your calculator to find  $AB$  where  $A = \begin{bmatrix} 1 & -1 & 1 \\ 0 & -2 & 1 \\ -2 & -3 & 0 \end{bmatrix}$  and  $B = \begin{bmatrix} 3 & -3 & 1 \\ -2 & 2 & -1 \\ -4 & 5 & -2 \end{bmatrix}$ .

**Example 4.11**

Which of the following is  $C^{-1}$ , where  $C = \begin{bmatrix} 5 & 3 \\ 2 & 1 \end{bmatrix}$ ?

a.  $A = \begin{bmatrix} \frac{1}{5} & \frac{1}{3} \\ \frac{1}{2} & 1 \end{bmatrix}$

b.  $B = \begin{bmatrix} -5 & -3 \\ -2 & -1 \end{bmatrix}$



d.  $D = \begin{bmatrix} -1 & 3 \\ 2 & -5 \end{bmatrix}$

**Example 4.12**

Determine – **by hand** – if they exist, the inverses of each of the following matrices.

$$A = \begin{bmatrix} -12 & -5 & -9 \\ -4 & -2 & -4 \\ -8 & -4 & -6 \end{bmatrix}, B = \begin{bmatrix} 1 & -1 & 0 \\ 1 & 0 & -1 \\ 6 & -2 & -3 \end{bmatrix} \text{ and } C = \begin{bmatrix} 7 & -12 \\ 18 & 4 \\ 0 & -9 \end{bmatrix}$$

**Example 4.13**

Determine a general formula for the inverse of the matrix  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$

**Example 4.14**

Let's use the formula derived in Example 4.13 to help us solve the system  $\begin{cases} 4x_1 + 3x_2 = 20 \\ -2x_1 + 5x_2 = -36 \end{cases}$ .

**Example 4.15**

Let  $A = \begin{bmatrix} 2 & 3 \\ 3 & 5 \end{bmatrix}$  and  $B = \begin{bmatrix} -4 & -1 \\ 7 & 2 \end{bmatrix}$ . Find, by hand,  $AB$ ,  $BA$ ,  $(AB)^T$ ,  $(BA)^T$ ,  $A^T B^T$ ,  $B^T A^T$ ,  $A^{-1}$ ,  $B^{-1}$ ,  $(A^{-1})^{-1}$ ,  $(AB)^{-1}$ ,  $(BA)^{-1}$ ,  $A^{-1} B^{-1}$ ,  $B^{-1} A^{-1}$ ,  $(A^{-1})^T$ , and  $(A^T)^{-1}$ . See what equals what.

**Definitions 4.1-4.5**

An  $m \times n$  **matrix** is a rectangular array of numbers with  $m$  rows and  $n$  columns.  $m$  and  $n$  are called the **dimensions** of the matrix.

Matrices are most commonly denoted by capital letters or single subscripted capital letters. The numbers (**entries**) of a matrix are denoted by double subscripted lower case letters.

When a matrix is explicitly written out, it is delineated by square brackets or parentheses. Abstractly, you will see things like  $A = [a_{ij}]$  where  $a_{ij}$  represents the entry in the  $i^{\text{th}}$  row and  $j^{\text{th}}$  column. When the number of rows or columns goes above 9, we separate the  $i$ s and  $j$ s by commas; we won't be going there.

A matrix with only one row is called a **row vector** and a matrix with only one column is called a **column vector**.

**Definitions 4.6 and 4.7**

**Matrix addition:** Two matrices with the exact same dimensions can be added or subtracted thusly:

$$\begin{bmatrix} a_{ij} \end{bmatrix} \pm \begin{bmatrix} b_{ij} \end{bmatrix} = \begin{bmatrix} a_{ij} \pm b_{ij} \end{bmatrix}$$

If either of the corresponding dimensions of two matrices is not the same, the matrices can neither be added nor subtracted.

**Scalar multiplication:** A matrix can be multiplied by a scalar (number) thusly:  $k \begin{bmatrix} a_{ij} \end{bmatrix} = \begin{bmatrix} k a_{ij} \end{bmatrix}$

**Definition 4.8**

The transpose of the  $m \times n$  matrix  $M$  is the  $n \times m$  matrix,  $M^T$ , that results from swapping the rows and columns of  $M$ .

**Definition 4.9: Generalized Matrix Multiplication**

If  $A$  is an  $m \times p$  matrix and  $B$  is a  $p \times n$  matrix, then the product  $AB$  is the  $m \times n$  matrix whose  $ij^{\text{th}}$  entry is the dot product of the  $i^{\text{th}}$  row of  $A$  and the  $j^{\text{th}}$  column of  $B$ .

If  $A$  is an  $m \times p$  matrix and  $B$  does not have  $p$  rows, then the product  $AB$  is not defined.

**Definitions 4.10-4.12 and Theorems 4.1-4.2**

The  $n \times n$  matrix with entries of 1 along the main diagonal and 0 in every other slot is called the  $n \times n$  identity matrix and is denoted as  $I_n$ .

If the products are defined,  $AI_n = A$  and/or  $I_n A = A$  for any compatible matrix  $A$ .

A square matrix  $A$  with dimensions  $n \times n$  is called invertible if there exists an  $n \times n$  matrix  $B$  with the property  $AB = I_n$ . If such a matrix exists we call it the inverse of  $A$  and symbolize it as  $A^{-1}$ .

If the  $n \times n$   $A$  is invertible, then  $AA^{-1} = A^{-1}A = I_n$ .

We do not define inverse matrices for non-square matrices.

**Algorithm 4.1**

A game plan for determining, **by hand**, the inverse of the matrix  $A = \begin{bmatrix} 1 & -1 & 0 \\ 1 & 0 & -1 \\ 6 & -2 & -3 \end{bmatrix}$  is shown below.

Let's execute the plan on the next page.

Let  $A^{-1} = \begin{bmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \\ b_{31} & b_{32} & b_{33} \end{bmatrix}$ . We need  $AA^{-1} = I_3$  which gives us:

$$\begin{bmatrix} 1 & -1 & 0 \\ 1 & 0 & -1 \\ 6 & -2 & -3 \end{bmatrix} \begin{bmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \\ b_{31} & b_{32} & b_{33} \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} b_{11} - b_{21} & b_{12} - b_{22} & b_{13} - b_{23} \\ b_{11} - b_{31} & b_{12} - b_{32} & b_{13} - b_{33} \\ 6b_{11} - 2b_{21} - 3b_{31} & 6b_{12} - 2b_{22} - 3b_{32} & 6b_{13} - 2b_{23} - 3b_{33} \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

If we equate the columns of the respective matrices, we come up with three systems of three equations with three unknowns. Specifically:

$$\begin{cases} b_{11} - b_{21} = 1 \\ b_{11} - b_{31} = 0 \\ 6b_{11} - 2b_{21} - 3b_{31} = 0 \end{cases}, \quad \begin{cases} b_{12} - b_{22} = 0 \\ b_{12} - b_{32} = 1 \\ 6b_{12} - 2b_{22} - 3b_{32} = 0 \end{cases}, \quad \text{and} \quad \begin{cases} b_{13} - b_{23} = 0 \\ b_{13} - b_{33} = 0 \\ 6b_{13} - 2b_{23} - 3b_{33} = 1 \end{cases}$$

A not-so-close inspection should convince you that not only are the coefficient matrices for all three of the systems identical, but they are all in fact the matrix  $A$ ! Since the row operations performed in Gaussian elimination are determined solely by the coefficient matrix, we may as well go ahead and solve all three systems simultaneously. The augmented matrix representation for these three systems is:

$$\left[ \begin{array}{ccc|ccc} 1 & -1 & 0 & 1 & 0 & 0 \\ 1 & 0 & -1 & 0 & 1 & 0 \\ 6 & -2 & -3 & 0 & 0 & 1 \end{array} \right]$$

common coefficient matrix

↑   ↑   ↑

Constant terms, from left to right, for the first, second, and third columns' related system of equations.