

$\begin{matrix} \nearrow \\ \text{\# of} \\ \text{rows} \end{matrix} \quad \begin{matrix} M \times n \\ \uparrow \\ \text{\# of columns} \end{matrix}$

A **column vector** is a matrix with only one column. Assuming that we limit the entries to real numbers, the set of all 2×1 (column) vectors is called \mathbb{R}^2 and the set of all 3×1 vectors is called \mathbb{R}^3 .

Scalar multiplication is the process of multiplying a vector by a real number; the process is effected by multiplying each entry in the vector by the scalar.

Vector addition and **vector subtraction** are effected by adding or subtracting the corresponding entries of the two vectors; both of these operations can only be performed between vectors with the same number of rows.

Example

Simplify $-2 \begin{bmatrix} -1 \\ -2 \end{bmatrix} + 3 \begin{bmatrix} 0 \\ -2 \end{bmatrix}$ and illustrate the process on Figure 1.

$$\begin{aligned}
 -2 \begin{bmatrix} -1 \\ -2 \end{bmatrix} + 3 \begin{bmatrix} 0 \\ -2 \end{bmatrix} &= \begin{bmatrix} 2 \\ 4 \end{bmatrix} + \begin{bmatrix} 0 \\ -6 \end{bmatrix} \\
 &= \begin{bmatrix} 2 \\ -2 \end{bmatrix}
 \end{aligned}$$

★ vector sum
 is $\begin{bmatrix} 2 \\ -2 \end{bmatrix}$




Figure 1: $-2 \begin{bmatrix} -1 \\ -2 \end{bmatrix} + 3 \begin{bmatrix} 0 \\ -2 \end{bmatrix}$

Suppose that $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p$ are all vectors in \mathbb{R}^n . We say that \vec{y} is a linear combination of $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p$ if and only if there exist scalars, c_1, c_2, \dots, c_p such that $\vec{y} = c_1 \vec{v}_1 + c_2 \vec{v}_2 + \dots + c_p \vec{v}_p$. If such scalars exist, they are called the weights in the linear combination.

Example

Let $\vec{a}_1 = \begin{bmatrix} 2 \\ -4 \end{bmatrix}$, $\vec{a}_2 = \begin{bmatrix} -5 \\ -1 \end{bmatrix}$, and $\vec{b} = \begin{bmatrix} 12 \\ -13 \end{bmatrix}$. Express \vec{b} as a linear combination of \vec{a}_1 and \vec{a}_2 .

Translation: Find numbers, c_1 and c_2 , such that

$$c_1 \begin{bmatrix} 2 \\ -4 \end{bmatrix} + c_2 \begin{bmatrix} -5 \\ -1 \end{bmatrix} = \begin{bmatrix} 12 \\ -13 \end{bmatrix}$$

System of equations

$$\begin{cases} 2c_1 - 5c_2 = 12 \\ -4c_1 - c_2 = -13 \end{cases}$$

$$\begin{bmatrix} 2c_1 \\ -4c_1 \end{bmatrix} + \begin{bmatrix} -5c_2 \\ -c_2 \end{bmatrix} = \begin{bmatrix} 12 \\ -13 \end{bmatrix}$$

$$\left[\begin{array}{cc|c} 2 & -5 & 12 \\ -4 & -1 & -13 \end{array} \right] \sim \left[\begin{array}{cc|c} 1 & 0 & 7/2 \\ 0 & 1 & -1 \end{array} \right] \therefore \begin{cases} c_1 = 7/2 \\ c_2 = -1 \end{cases}$$

$$\therefore \begin{bmatrix} 12 \\ -13 \end{bmatrix} = \frac{7}{2} \begin{bmatrix} 2 \\ -4 \end{bmatrix} + (-1) \begin{bmatrix} -5 \\ -1 \end{bmatrix}$$

Vocab: The weight on \vec{a}_1 is $\frac{7}{2}$ and the weight on \vec{a}_2 is -1

Suppose that $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p$ are all vectors in \mathbb{R}^n . The set of all linear combinations of $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p$ is called the span of $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p$.

General Solution:
$$\begin{cases} x_1 = -2x_3 + 14/11 \\ x_2 = -x_3 + 9/11 \\ x_3 = \text{free} \end{cases}$$
 Letting $x_3 = 0$, we see that

MTH 261 - Mr. Simonds' class

Example Hence let's use x_i as the i^{th} weight. $\vec{b} = \frac{14}{11}\vec{c}_1 + \frac{9}{11}\vec{c}_2 + 0\vec{c}_3$

Let $A = \begin{bmatrix} 3 & -1 & 5 \\ -2 & 0 & -4 \\ -1 & 4 & 2 \end{bmatrix}$ and $\vec{b} = \begin{bmatrix} 3 \\ h \\ 2 \end{bmatrix}$. Find the value of h if \vec{b} is in the span of the columns of A .

Translation 1: We want \vec{b} to be expressed as a linear combination of the columns of A .

Translation 2: $x_1 \begin{bmatrix} 3 \\ -2 \\ -1 \end{bmatrix} + x_2 \begin{bmatrix} -1 \\ 0 \\ 4 \end{bmatrix} + x_3 \begin{bmatrix} 5 \\ -4 \\ 2 \end{bmatrix} = \begin{bmatrix} 3 \\ h \\ 2 \end{bmatrix}$

We want this to have a solution.

System of equations:
$$\begin{cases} 3x_1 - x_2 + 5x_3 = 3 \\ -2x_1 - 4x_3 = h \\ -x_1 + 4x_2 + 2x_3 = 2 \end{cases}$$

$$\left[\begin{array}{ccc|c} 3 & -1 & 5 & 3 \\ -2 & 0 & -4 & h \\ -1 & 4 & 2 & 2 \end{array} \right] R_1 \leftrightarrow R_3 \left[\begin{array}{ccc|c} -1 & 4 & 2 & 2 \\ -2 & 0 & -4 & h \\ 3 & -1 & 5 & 3 \end{array} \right]$$


$$\begin{aligned} -2R_1 + R_2 &\rightarrow R_2 \\ 3R_1 + R_3 &\rightarrow R_3 \end{aligned} \left[\begin{array}{ccc|c} -1 & 4 & 2 & 2 \\ 0 & -8 & -8 & -4+h \\ 0 & 11 & 11 & 9 \end{array} \right]$$

$$\frac{11}{8}R_2 + R_3 \rightarrow R_3 \left[\begin{array}{ccc|c} -1 & 4 & 2 & 2 \\ 0 & -8 & -8 & -4+h \\ 0 & 0 & 0 & \frac{7}{2} + \frac{11}{8}h \end{array} \right]$$

The system will have a solution iff $\frac{7}{2} + \frac{11}{8}h = 0$.

$\therefore \vec{b}$ is in the span of the columns of A iff $h = -\frac{28}{11}$

Check
$$\left[\begin{array}{ccc|c} 3 & -1 & 5 & 3 \\ -2 & 0 & -4 & -28/11 \\ -1 & 4 & 2 & 2 \end{array} \right] \sim \left[\begin{array}{ccc|c} 1 & 0 & 2 & 14/11 \\ 0 & 1 & 1 & 9/11 \\ 0 & 0 & 0 & 0 \end{array} \right] \checkmark$$



MTH 261 – Mr. Simonds' class

$$\text{Balance : } m_1 d_1 + m_2 d_2 = 0$$

Application: Center of Mass

The center of mass (or center of gravity) of an object is the average of the product of mass points in the object with their relative distance from a fixed reference point. The concept is most easily understood if one thinks about a teeter-totter. When two people of different mass sit on a teeter totter of uniform density, the center of mass is clearly going to be closer to the heavier person than the lighter person; this phenomenon is reflected in the fact that if the teeter-totter is to stay in balance, the heavier person needs to sit closer to the axis-point than the lighter person. Unfortunately, we can't define the center of mass in all cases as the balance point, because the center of mass is frequently not even on the object! This is easy to see if you think about washer (as in nuts and bolts) of uniform density; the center of mass is clearly the center point of the hole in the middle of the washer.

For most objects, it takes a double or triple integral to calculate the center of mass. In some simple situations, however, the point can be determined by a simple formula. For example, if we have a triangular lamina referenced to the xy -plane, the center of mass (\bar{v}) can be determined using the

formula $\bar{v} = \frac{1}{3}[\bar{v}_1 + \bar{v}_2 + \bar{v}_3]$ where \bar{v}_1 , \bar{v}_2 , and \bar{v}_3 represent the vectors from the origin to the three vertices of the triangle; for ease of reference, we shall contextually refer to these vectors as points.

That formula is a special case of the more general formula for the center of mass of several mass points.

The center of mass (\bar{v}) of n mass points is: $\bar{v} = \frac{1}{m} \sum_{i=1}^n [m_i v_i]$ where m_i is the mass at point v_i and m is the sum of all the masses.

Example

Find the center of mass of the triangular lamina outlined in Figure 2 assuming that the lamina has uniform density.




Figure 2: lamina of uniform density and thickness

The center of mass
(assuming uniform density) is

$$\frac{1}{3} \left(\begin{bmatrix} -3 \\ -4 \end{bmatrix} + \begin{bmatrix} 4 \\ 5 \end{bmatrix} + \begin{bmatrix} 5 \\ -4 \end{bmatrix} \right) = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$$

Example

In the last example we found the center of mass using the formula $\vec{v} = \frac{1}{m} \sum_{i=1}^n [m_i, v_i]$ and assuming that there was 1 g of mass at each of the vertices. Suppose that we had 9 additional grams of mass to distribute among the vertices. How should the mass be distributed so that the center of mass of the lamina shifts to the point $(3, -2)$?

$$C \text{ of } m = \begin{bmatrix} 3 \\ -2 \end{bmatrix}$$

$$\frac{1}{12} \left(x_1 \begin{bmatrix} -3 \\ -4 \end{bmatrix} + x_2 \begin{bmatrix} 4 \\ 5 \end{bmatrix} + x_3 \begin{bmatrix} 5 \\ -4 \end{bmatrix} \right) = \begin{bmatrix} 3 \\ -2 \end{bmatrix}$$

$$\text{also } x_1 + x_2 + x_3 = 12$$

$$\text{System: } \begin{cases} -3x_1 + 4x_2 + 5x_3 = 36 \\ -4x_1 + 5x_2 - 4x_3 = -24 \\ x_1 + x_2 + x_3 = 12 \end{cases}$$

$$\left[\begin{array}{ccc|c} -3 & 4 & 5 & 36 \\ -4 & 5 & -4 & -24 \\ 1 & 1 & 1 & 12 \end{array} \right] \sim \left[\begin{array}{ccc|c} 1 & 0 & 0 & 8/3 \\ 0 & 1 & 0 & 8/3 \\ 0 & 0 & 1 & 20/3 \end{array} \right]$$




Figure 2: lamina of uniform density and thickness

$$\begin{aligned} x_1 &= \text{total mass distributed to A} \\ x_2 &= \text{total mass distributed to B} \\ x_3 &= \text{total mass distributed to C} \end{aligned}$$

Vector Equations – Section 1.3 / Matrix Equations – Section 1.4 | 5

\therefore we need to distribute $5/3$ g to each of the pairs A + B and the remaining $17/3$ g to point C.

Definition

The product of an $m \times n$ matrix, A , and $n \times 1$ vector, \vec{x} , is defined by $A\vec{x} = \sum_{i=1}^n x_i \vec{a}_i$ where x_i is the entry in the i^{th} row of \vec{x} and \vec{a}_i is the i^{th} column of A (treating the columns of A as vectors). You cannot find the product $A\vec{x}$ unless the number of columns of A is equal to the number of rows of \vec{x} .

Example

Let $A = \begin{bmatrix} 2 & -6 & -1 \\ 0 & 5 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 1 & -3 \\ 4 & 3 \\ -2 & 7 \end{bmatrix}$, $\vec{u} = \begin{bmatrix} 10 \\ 2 \end{bmatrix}$, and $\vec{v} = \begin{bmatrix} -2 \\ 4 \\ 8 \end{bmatrix}$. Find each of the following products (where possible): $A\vec{u}$, $A\vec{v}$, $B\vec{u}$, and $B\vec{v}$.

inside dimensions are unequal

2×3 2×1

$A\vec{u}$ cannot be executed

$$2 \times 3 \quad 3 \times 1 \quad A\vec{v} = \begin{bmatrix} 2 & -6 & -1 \\ 0 & 5 & 2 \end{bmatrix} \begin{bmatrix} -2 \\ 4 \\ 8 \end{bmatrix}$$

$$= -2 \begin{bmatrix} 2 \\ 0 \end{bmatrix} + 4 \begin{bmatrix} -6 \\ 5 \end{bmatrix} + 8 \begin{bmatrix} -1 \\ 2 \end{bmatrix}$$

$$= \begin{bmatrix} (-2)(2) + (4)(-6) + (8)(-1) \\ (-2)(0) + (4)(5) + (8)(2) \end{bmatrix}$$

$$= \begin{bmatrix} -36 \\ 36 \end{bmatrix}$$

$B\vec{v}$ is a no-go 3×2 2×1 mismatch
 3×2 2×1 result is 3×1

$$B\vec{u} = \begin{bmatrix} 1 & -3 \\ 4 & 3 \\ -2 & 7 \end{bmatrix} \begin{bmatrix} 10 \\ 2 \end{bmatrix}$$

$$= 10 \begin{bmatrix} 1 \\ 4 \\ -2 \end{bmatrix} + 2 \begin{bmatrix} -3 \\ 3 \\ 7 \end{bmatrix}$$

$$= \begin{bmatrix} 4 \\ 46 \\ -6 \end{bmatrix}$$

Example

Write $\begin{bmatrix} -1 & 2 & 5 \\ 8 & -2 & 0 \\ 1 & -3 & 6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 7 \\ 11 \\ -3 \end{bmatrix}$ as a vector equation.

vector eq. $x_1 \begin{bmatrix} -1 \\ 8 \\ 1 \end{bmatrix} + x_2 \begin{bmatrix} 2 \\ -2 \\ -3 \end{bmatrix} + x_3 \begin{bmatrix} 5 \\ 0 \\ 6 \end{bmatrix} = \begin{bmatrix} 7 \\ 11 \\ -3 \end{bmatrix}$

System of equations $\begin{cases} -x_1 + 2x_2 + 5x_3 = 7 \\ 8x_1 - 2x_2 = 11 \\ x_1 - 3x_2 + 6x_3 = -3 \end{cases}$

Example

Write the system $\begin{cases} 2x_1 - 3x_2 + x_3 - 2x_4 = 0 \\ 5x_1 - x_2 + x_4 = 9 \end{cases}$ as a matrix equation of form $A\vec{x} = \vec{b}$.

$$\begin{bmatrix} 2 & -3 & 1 & -2 \\ 5 & -1 & 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 0 \\ 9 \end{bmatrix}$$

Coefficient matrix

$$\vec{b} = \{ \vec{v}_1, \vec{v}_2 \}$$

The equation $A\vec{x} = \vec{b}$ has a solution if and only if \vec{b} can be expressed as a linear combination of the columns of A ; that is, $A\vec{x} = \vec{b}$ has a solution if and only if \vec{b} is in the span of the columns of A .

Later in the term the span of the columns of A will be defined as the column space of A . It follows that the equation $A\vec{x} = \vec{b}$ has a solution if and only if \vec{b} is in the column space of A .

Example

Describe the column spaces of the matrixes A , B , and C where $A = \begin{bmatrix} -2 \\ 5 \end{bmatrix}$, $B = \begin{bmatrix} 0 & 0 \\ 2 & -1 \\ 3 & -4 \end{bmatrix}$, and

$$C = \begin{bmatrix} -1 & 2 \\ 3 & -4 \\ 6 & 7 \\ -1 & 1 \end{bmatrix}.$$

A: The column space is the set of vectors $\left\{ x_1 \begin{bmatrix} -2 \\ 5 \end{bmatrix} \mid x_1 \in \mathbb{R} \right\}$
 This is a line through the origin parallel to $\begin{bmatrix} -2 \\ 5 \end{bmatrix}$.

$$B: \left\{ x_1 \begin{bmatrix} 0 \\ 2 \\ 3 \end{bmatrix} + x_2 \begin{bmatrix} 0 \\ -1 \\ -4 \end{bmatrix} \mid x_1, x_2 \in \mathbb{R} \right\}$$

Two non-multiple vectors always span a plane.
 linear independent vectors

In this case, that plane is the x_2x_3 -plane.

$$C: \left\{ x_1 \begin{bmatrix} -1 \\ 3 \\ 6 \\ -1 \end{bmatrix} + x_2 \begin{bmatrix} 2 \\ -4 \\ 7 \\ 1 \end{bmatrix} \mid x_1, x_2 \in \mathbb{R} \right\}$$

Even though this space comes from the non-singular 4th dimension, this span is a plane.