

Solutions to “Some Additional Practice for the Midterm Exam”

1. a. Convert 24° into radians.

$$\begin{aligned} 24^\circ &= 24^\circ \cdot \frac{2\pi \text{ rad.}}{360^\circ} \\ &= \frac{2\pi}{15} \text{ rad.} \end{aligned}$$

- b. Convert $\frac{3}{2}$ radians into degrees.

$$\begin{aligned} \frac{3}{2} \text{ rad.} &= \frac{3}{2} \text{ rad.} \cdot \frac{360^\circ}{2\pi \text{ rad.}} \\ &= \frac{540^\circ}{2\pi} \\ &= \frac{270^\circ}{\pi} \end{aligned}$$

2. Find the arc-length spanned by an angle measuring 24° on a circle of radius 30 feet.

In part a of #1 we found that $24^\circ = \frac{2\pi}{15}$ rad. Thus,

$$\begin{aligned} s &= r\theta \\ &= 30 \cdot \frac{2\pi}{15} \\ &= 4\pi \end{aligned}$$

and we see that the arc-length spanned by an angle measuring 24° on a circle of radius 30 feet is 4π feet.

3. Evaluate the following expressions:

a. $\sin(225^\circ) = -\sin(45^\circ)$ (since 45° is the reference angle for 225° and sine is negative in the 3rd quadrant)

$$= -\frac{\sqrt{2}}{2}$$

- b. $\sin\left(\frac{5\pi}{3}\right) = -\sin\left(\frac{\pi}{3}\right)$ (since $\frac{\pi}{3}$ is the reference angle for $\frac{5\pi}{3}$
and sine is negative in the 4th quadrant)
 $= -\frac{\sqrt{3}}{2}$
- c. $\cos(300^\circ) = \cos(60^\circ)$ (since 60° is the reference angle for 300°)
 $= \frac{1}{2}$
- d. $\cos\left(\frac{17\pi}{6}\right) = \cos\left(\frac{5\pi}{6}\right)$ (since $\frac{5\pi}{6}$ is coterminal with $\frac{17\pi}{6}$)
 $= -\cos\left(\frac{\pi}{6}\right)$ (since $\frac{\pi}{6}$ is the reference angle for $\frac{5\pi}{6}$
and cosine is negative in the 2nd quadrant)
 $= -\frac{\sqrt{3}}{2}$
- e. $\tan\left(\frac{3\pi}{4}\right) = \frac{\sin\left(\frac{3\pi}{4}\right)}{\cos\left(\frac{3\pi}{4}\right)}$
 $= \frac{\sin\left(\frac{\pi}{4}\right)}{-\cos\left(\frac{\pi}{4}\right)}$ (since $\frac{\pi}{4}$ is the reference angle for $\frac{3\pi}{4}$
and cosine is negative in the 2nd quadrant)
 $= \frac{\frac{\sqrt{2}}{2}}{-\frac{\sqrt{2}}{2}}$
 $= -1$
- f. $\sec(3600^\circ) = \frac{1}{\cos(3600^\circ)}$
 $= \frac{1}{\cos(360^\circ)}$ (since 360° is coterminal with 3600°)
 $= \frac{1}{1}$
 $= 1$

4. If $\sin(\theta) = \frac{3}{4}$ and θ is in the second quadrant, find the exact value of the following expressions.

a. $\cos(\theta)$

We can use the Pythagorean identity to find $\cos(\theta)$:

$$\begin{aligned} \sin^2(\theta) + \cos^2(\theta) &= 1 \\ \Rightarrow \left(\frac{3}{4}\right)^2 + \cos^2(\theta) &= 1 \\ \Rightarrow \cos^2(\theta) &= 1 - \frac{9}{16} \\ \Rightarrow \cos^2(\theta) &= \frac{7}{16} \\ \Rightarrow \cos(\theta) &= -\frac{\sqrt{7}}{4} \quad (\text{note that we take the negative square root} \\ &\quad \text{since cosine is negative in the second quadrant}) \end{aligned}$$

b. $\sin(\theta + 2\pi) = \sin(\theta)$ (since the period of $y = \sin(\theta)$ is 2π)

$$= \frac{3}{4}$$

c. $\sin(-\theta) = -\sin(\theta)$ (due to the symmetry of a circle and the definition of sine)

$$= -\frac{3}{4}$$

d. $\tan(\theta) = \frac{\sin(\theta)}{\cos(\theta)}$

$$\begin{aligned} &= \frac{\frac{3}{4}}{-\frac{\sqrt{7}}{4}} \\ &= -\frac{3}{\sqrt{7}} \end{aligned}$$

e. $\csc(\theta) = \frac{1}{\sin(\theta)}$

$$\begin{aligned} &= \frac{1}{\frac{3}{4}} \\ &= \frac{4}{3} \end{aligned}$$

$$\begin{aligned}
 \text{f. } \sec(\theta) &= \frac{1}{\cos(\theta)} \\
 &= \frac{1}{-\frac{\sqrt{7}}{4}} \\
 &= -\frac{4}{\sqrt{7}}
 \end{aligned}$$

5. Find all of the solutions to the following equations on the interval $[0, 2\pi)$:

a. $2\cos(\theta) = 1$

$$\begin{aligned}
 2\cos(\theta) &= 1 \\
 \Rightarrow \cos(\theta) &= \frac{1}{2}
 \end{aligned}$$

In the first quadrant, we know that $\theta = \frac{\pi}{3}$ is a solution.

In the second and third quadrant cosine is negative, so there are no solutions.

In the fourth quadrant we know that $\theta = \frac{5\pi}{3}$ is a solution.

Thus the solution set on the interval $[0, 2\pi)$ is $\left\{\frac{\pi}{3}, \frac{5\pi}{3}\right\}$.

b. $\sin(2\theta) + 3 = 4$

$$\begin{aligned}
 \sin(2\theta) + 3 &= 4 \\
 \Rightarrow \sin(2\theta) &= 1
 \end{aligned}$$

Since $\sin(x) = 1$ when $x = \frac{\pi}{2} + 2k\pi$ (where $k \in \mathbb{Z}$), we need:

$$\begin{aligned}
 2\theta &= \frac{\pi}{2} + 2k\pi \\
 \Rightarrow \theta &= \frac{\frac{\pi}{2} + 2k\pi}{2} \\
 \Rightarrow \theta &= \frac{\pi}{4} + k\pi
 \end{aligned}$$

Since $0 \leq \theta < 2\pi$, we can only take $k = 0$ and $k = 1$ in the statement

above. Thus, the solution set is $\left\{\frac{\pi}{4}, \frac{5\pi}{4}\right\}$.

6. Find *all* of the solutions to the following equations:

a. $3\sin(x) + 4 = 5$

$$\begin{aligned} 3\sin(x) + 4 &= 5 \\ \Rightarrow 3\sin(x) &= 1 \\ \Rightarrow \sin(x) &= \frac{1}{3} \end{aligned}$$

Since this sine value isn't one we are familiar with, we will need to use the inverse sine function (i.e., arcsine function).

$$\begin{aligned} \sin(x) &= \frac{1}{3} \\ \Rightarrow x &= \sin^{-1}\left(\frac{1}{3}\right) + 2k\pi, \\ \text{or } x &= \pi - \sin^{-1}\left(\frac{1}{3}\right) + 2k\pi \quad \text{where } k \in \mathbb{Z} \end{aligned}$$

b. $7 + 3\sqrt{2}\cos(4t) = 4$

$$\begin{aligned} 7 + 3\sqrt{2}\cos(4t) &= 4 \\ \Rightarrow 3\sqrt{2}\cos(4t) &= -3 \\ \Rightarrow \cos(4t) &= \frac{-3}{3\sqrt{2}} \\ \Rightarrow \cos(4t) &= -\frac{1}{\sqrt{2}} \\ \Rightarrow \left\{ \begin{array}{l} 4t = \frac{3\pi}{4} + 2k\pi, \quad k \in \mathbb{Z} \\ \text{OR} \\ 4t = -\frac{3\pi}{4} + 2k\pi, \quad k \in \mathbb{Z} \end{array} \right. \\ \Rightarrow \left\{ \begin{array}{l} t = \frac{3\pi}{16} + \frac{k\pi}{2}, \quad k \in \mathbb{Z} \\ \text{OR} \\ t = -\frac{3\pi}{16} + \frac{k\pi}{2}, \quad k \in \mathbb{Z} \end{array} \right. \end{aligned}$$

7. Use the sine and cosine functions to find the coordinates of the point P in Figure 1 that is specified by $\frac{7\pi}{6}$ on the circumference of a circle of radius 3 units.




Figure 1

Coordinates of P :

$$\begin{aligned} \left(3 \cos\left(\frac{7\pi}{6}\right), 3 \sin\left(\frac{7\pi}{6}\right)\right) &= \left(3 \cdot \left(-\frac{\sqrt{3}}{2}\right), 3 \cdot \left(-\frac{1}{2}\right)\right) \\ &= \left(-\frac{3\sqrt{3}}{2}, -\frac{3}{2}\right) \end{aligned}$$


8. Sketch a graph of $g(t) = 3 \sin\left(2t + \frac{\pi}{4}\right) - 2$. State the period, midline, and amplitude.

First, let's write the function in "standard form":

$$\begin{aligned} g(t) &= 3 \sin\left(2t + \frac{\pi}{4}\right) - 2 \\ &= 3 \sin\left(2\left(t + \frac{\pi}{8}\right)\right) - 2 \end{aligned}$$

Thus, this is a function of the form $g(t) = A \sin(\omega(t - h)) + k$ where $A = 3$, $\omega = 2$, $h = -\frac{\pi}{8}$, and $k = -2$. Since $A = 3$ the function has **amplitude 3 units**. Using the fact that $\omega = 2$, we can find the period: $\text{period} = \frac{2\pi}{\omega} = \frac{2\pi}{2} = \pi$.

So the **period is π units**. Since $k = -2$, the **midline is $y = -2$** . Since $h = -\frac{\pi}{8}$, we need to "start" our sine wave at $t = -\frac{\pi}{8}$, i.e., shift the wave left $\frac{\pi}{8}$ units. Below is a graph of $g(t) = 3 \sin\left(2\left(t + \frac{\pi}{8}\right)\right) - 2$.



A graph of $g(t) = 3 \sin\left(2t + \frac{\pi}{4}\right) - 2 = 3 \sin\left(2\left(t + \frac{\pi}{8}\right)\right) - 2$.

9. Find four algebraic rules (one using positive sine, one using negative sine, one using positive cosine, and one using negative cosine) for the function $y = f(t)$ graphed in Figure 2.




Figure 2: Graph of $y = f(t)$.

Whether we use sine or cosine for our rule, it will have form $f(t) = \pm|A|\sin(\omega(t - h)) + k$ or $f(t) = \pm|A|\cos(\omega(t - h)) + k$ so we can start by determining the values of $|A|$, ω , and k .

- the amplitude is 4 units so we know that $|A| = 4$
- the midline is $y = 8$ so we know that $k = 8$
- the period is 1 unit so we know that ω must satisfy $1 = 2\pi \cdot \frac{1}{\omega}$. Thus,

$$1 = 2\pi \cdot \frac{1}{\omega}$$

$$\Rightarrow \omega = \frac{2\pi}{1} = 2\pi$$

Since a positive sine wave “starts” at $t = 0.5$, we can use $h = 0.5$ to construct a rule:

$$f(t) = 4\sin(2\pi(t - 0.5)) + 8$$

Since a negative (reflected) sine wave “starts” at $t = 0$, we can use $h = 0$:

$$f(t) = -4\sin(2\pi \cdot t) + 8$$

Since a positive cosine wave “starts” at $t = -0.25$, we can use $h = -0.25$:

$$f(t) = 4\cos(2\pi(t + 0.25)) + 8$$

Since a negative (reflected) cosine wave “starts” at $t = 0.25$, we can use $h = 0.25$:

$$f(t) = -4\cos(2\pi(t - 0.25)) + 8$$

(Note that there are many other possible answers.)

b. Use one of your answer to part a to find exact solutions to $f(t) = 10$.

$$\begin{aligned}
 f(t) &= 10 \\
 \Rightarrow -4\sin(2\pi t) + 8 &= 10 \\
 \Rightarrow -4\sin(2\pi t) &= 2 \\
 \Rightarrow \sin(2\pi t) &= -\frac{1}{2} \\
 \Rightarrow 2\pi t = -\frac{\pi}{6} + 2k\pi &\quad \text{or} \quad 2\pi t = \frac{7\pi}{6} + 2k\pi, \quad k \in \mathbb{Z} \\
 \Rightarrow t = \frac{-\frac{\pi}{6} + 2k\pi}{2\pi} &\quad \text{or} \quad t = \frac{\frac{7\pi}{6} + 2k\pi}{2\pi}, \quad k \in \mathbb{Z} \\
 \Rightarrow t = -\frac{1}{12} + k &\quad \text{or} \quad t = \frac{7}{12} + k, \quad k \in \mathbb{Z}
 \end{aligned}$$

10. Evaluate the following:

a. $\sin\left(\cos^{-1}\left(-\frac{1}{2}\right)\right)$

$$\begin{aligned}
 \sin\left(\cos^{-1}\left(-\frac{1}{2}\right)\right) &= \sin\left(\frac{2\pi}{3}\right) \quad (\text{since the range of } y = \cos^{-1}(x) \text{ is } [0, \pi]) \\
 &= \frac{\sqrt{3}}{2}
 \end{aligned}$$

b. $\sin^{-1}\left(\sin\left(\frac{7\pi}{4}\right)\right)$

$$\begin{aligned}
 \sin^{-1}\left(\sin\left(\frac{7\pi}{4}\right)\right) &= \sin^{-1}\left(-\frac{\sqrt{2}}{2}\right) \\
 &= -\frac{\pi}{4} \quad (\text{since the range of } y = \sin^{-1}(x) \text{ is } \left[-\frac{\pi}{2}, \frac{\pi}{2}\right])
 \end{aligned}$$

c. $\cos^{-1}\left(\sin\left(\frac{4\pi}{3}\right)\right)$

$$\begin{aligned}
 \cos^{-1}\left(\sin\left(\frac{4\pi}{3}\right)\right) &= \cos^{-1}\left(-\frac{\sqrt{3}}{2}\right) \\
 &= \frac{5\pi}{6} \quad (\text{since the range of } y = \cos^{-1}(x) \text{ is } [0, \pi])
 \end{aligned}$$